

# Knots and Random Walks in Vibrated Granular Chains

Eli Ben-Naim

*Los Alamos National Laboratory*

Zahir Daya (Los Alamos)

Aaron Lauda (UC Riverside)

Peter Vorobieff (New Mexico)

Robert Ecke (Los Alamos)



Phys. Rev. Lett. **86**, 1414 (2001)

# Plan

I Knots






II Vibrated Knot Experiment

III Diffusion Theory

IV Experiment vs Theory

V Conclusions & Outlook

# Knots in Physical Systems



Knots in DNA strands

Tying a microtubule with optical tweezers

Knotted jets in accretion disks (MHD)

Strain on knot (MD)

Wang JMB 71

Itoh, Nature 99

F Thomsen 99

Wasserman, Nature 99

# Knots & Topological Constraints

- Knots happen

Whittington JCP 88

$$\text{probability(no knot)} \sim \exp(-N/N_0)$$

- Knots tighten ( $T = \infty$ )

Sommer JPA 92

$$n/N \rightarrow 0 \quad \text{when} \quad N \rightarrow \infty$$

- Reduce size of chain ( $m = \text{knot complexity}$ )

$$R \sim N^\nu m^{-\alpha} \quad \alpha = \nu - 1/3$$

- Reduce accessible phase space

- Large relaxation times

de Gennes, Edwards

$$\tau_{\text{reptation}} \sim N^3$$

- Weaken macromolecule

- Bio: affect chemistry, function

# Granular Chains

## Mechanical analog of bead-spring model


$$U(\{\mathbf{R}_i\}) = v_0 \sum_{i \neq j} \delta(\mathbf{R}_i - \mathbf{R}_j) + \frac{3}{2b^2} \sum_i (\mathbf{R}_i - \mathbf{R}_{i+1})^2$$

- Beads/rods interact via hard core repulsions
- Rods act as springs (nonlinear, dissipative)
- Inelastic collisions: bead-bead, bead-plate
- Vibrating plate supplies energy
- **Athermal, nonequilibrium driving**

## Advantages

- Number of beads can be controlled
- Topological constraints: can be prepared, observed directly

# Vibrated Knot Experiment




- $t = 0$ : trefoil knot placed at chain center
- Parameters
  - Number of monomers:  $30 < N < 270$
  - Minimal knot size:  $N_0 = 15$
- Driving conditions
  - Frequency:  $\nu = 13Hz$
  - Acceleration:  $\Gamma = A\omega^2/g = 3.4$

**Only measurement: opening time  $t$**

1. Average opening time  $\tau(N)$ ?
2. Survival probability  $S(t, N)$ ?  
Distribution of opening times  $R(t, N)$ ?

# The Average Opening Time



Average over 400 independent measurements

$$\tau(N) \sim (N - N_0)^\nu \quad \nu = 2.0 \pm 0.1$$

**Opening time is diffusive**


# The Survival Probability

- $S(t, N)$  Probability knot “alive” at time  $t$
- $R(t, N)$  Probability knot opens at time  $t$

$$S(t, N) = 1 - \int_0^t dt' R(t', N)$$

- $S(t, N)$  obeys scaling


$$S(t, N) = F(z) \quad z = \frac{t}{\tau(N)}$$



**$\tau$  only relevant time scale**



# Theoretical Model



## Assumptions

- Knot  $\equiv$  3 exclusion points
- Points hop randomly
- Points move independently (no correlation)
- Points are equivalent (size =  $N_0/3$ )

## 3 Random Walk Model

- 1D walks with excluded volume interaction
- first point reaches boundary  $\rightarrow$  knot opens

# Diffusion in 3D

$$1 < x_1 < x_2 < x_3 < N - N_0 \quad \longrightarrow \quad 0 < x < y < z < 1$$

$$\frac{\partial}{\partial t} P(x, y, z, t) = \nabla^2 P(x, y, z, t)$$

- Boundary conditions

Absorbing:  $P|_{x=0} = P|_{z=1} = 0$

Reflecting:  $(\partial_x - \partial_y)P|_{x=y} = (\partial_y - \partial_z)P|_{y=z} = 0$

- Initial conditions  $P|_{t=0} = \delta(x-x_0)\delta(y-x_0)\delta(z-x_0)$

- Survival probability

$$S_3(t) = \int_0^1 dx \int_x^1 dy \int_y^1 dz P(x, y, z, t)$$

**3 walks in 1D  $\equiv$  1 walk in 3D**

# Product Solution

- Product of 1D solutions

$$P(x, y, z, t) = 3! p(x, t)p(y, t)p(z, t)$$

- 1D case  $p|_{x=0} = p|_{x=1} = 0$   $p|_{t=0} = \delta(x - x_0)$

$$p_t(x, t) = p_{xx}(x, t)$$

- 1 walk survival probability  $s(t) = \int_0^1 dx p(x, t)$

$$s(t) = \frac{4}{\pi} \sum_{k=0}^{\infty} \frac{\sin[(2k+1)\pi x_0]}{2k+1} e^{-(2k+1)^2 \pi^2 t}$$

- $m$  interacting walks survival probability


$$S_m(t) = [s(t)]^m$$

- Average opening time

$$\langle t \rangle \simeq \tau_m \frac{(N - N_0)^2}{D} \quad \tau_3 = 0.056213$$

**Reduced to noninteracting problem**

## Alternative Derivation




Exchange identities of walkers when paths cross

# Experiment vs. Theory

- Work with scaling variable  $z = t/\tau$  ( $\langle z \rangle = 1$ )
- Combine different data sets (6000 pts)
- Fluctuations  $\sigma^2 = \langle z^2 \rangle - \langle z \rangle^2$

$$\sigma_{\text{exp}} = 0.62(1), \quad \sigma_{\text{theory}} = 0.63047 \quad (< 2\%)$$



No fitting parameters!


**Excellent quantitative agreement**

# The Exit Time Probability

## Scaling function

$$R(t, N) = \frac{1}{\tau} G(z) \quad z = \frac{t}{\tau(N)}$$

$$G(z) = -\frac{d}{dz} F(z)$$




# Large Exit Times

- Largest decay time dominates
- Large time tail is exponentially small

$$F(z) \sim e^{-\beta z} \quad z \gg 1$$

- Decay coefficient  $\beta = m\pi^2\tau_m$

$$\beta_{\text{exp}} = 1.65(2) \quad \beta_{\text{theory}} = 1.66440 \quad (1\%)$$




# Small Exit Times

- Exponentially small (in  $1/z$ ) tail

$$1 - F(z) \sim z^{1/2} e^{-\alpha/z} \quad z \ll 1$$

- Decay coefficient  $\alpha = 1/16\tau_m$

$$\alpha_{\text{exp}} = 1.2(1) \quad \alpha_{\text{theory}} = 1.11184 \quad (10\%)$$



**Larger discrepancy**



# Heuristic Argument (short times)

- Use scaling form

$$S(t, N) \sim F\left(\frac{t}{N^2}\right)$$

- Smallest exit time  $t = \frac{N}{2}$ ,  $1 - S \sim 2^{-N/2}$

$$1 - F\left(\frac{2}{N}\right) \sim e^{-\alpha N} \quad N \rightarrow \infty$$

$$1 - F(z) \sim e^{-\alpha/z} \quad z \rightarrow 0$$





- Analytic calculation: Laplace transform of  $s(t)$  + steepest descent

$$1 - F(z) \sim z^{1/2} e^{-\alpha/z}$$


- Complex knots:  $e^{-1/t} \sim m^{-1}$

$$\tau \sim \sigma \sim \frac{1}{\ln m} \quad m \gg 1$$

# Different knots ( $m = 1, 3, 5, 7$ )



# Off-Center Initial Conditions



Decay coefficient independent of  $x_0$

$$S_m(t) \simeq A(x_0)e^{-m\pi^2 t}$$

**Eventually, initial conditions are forgotten**

# Knots Opening & the Gambler Ruin Problem

- The exit probability

$$\nabla^2 E(x_1, \dots, x_d) = 0$$

- Linear in 1D:  $E(x_0) = x_0$
- In general dimension  $d \equiv m$

$$E(x_0) \sim (x_0)^d \quad x_0 \ll 1$$

- The average exit time

$$D\nabla^2 T(x_1, \dots, x_d) = -1$$



- General solution

$$E(x_0) = \frac{d}{2} \left(\frac{4}{\pi}\right)^d \sum_{k_1=1}^{\infty} \sum_{k_2=0}^{\infty} \cdots \sum_{k_d=0}^{\infty} \frac{(-1)^{k_1-1} k_1 \sin[k_1 \pi x_0]}{k_1^2 + \sum_{i=2}^d (2k_i + 1)^2} \prod_{i=2}^d \frac{\sin[(2k_i + 1)\pi x_0]}{(2k_i + 1)}$$

$$T(x_0) = \frac{1}{\pi^2} \left(\frac{4}{\pi}\right)^d \sum_{k_1=0}^{\infty} \cdots \sum_{k_d=0}^{\infty} \frac{1}{\sum_{i=1}^d (2k_i + 1)^2} \prod_{i=1}^d \frac{\sin[(2k_i + 1)\pi x_0]}{(2k_i + 1)}$$

**Knot opening  $\equiv$  3 gamblers ruin problem with fixed wealth hierarchy**

# Predictions



- Good agreement for  $S(t)$ ,  $S_{\text{far}}(t)$ ,  $S_{\text{close}}(t)$
- Poor agreement for  $E(x_0)$ ,  $T(x_0)$
- Current data insufficient (600pts)

**Fluctuations diverge near boundary**

# Conclusions

- Knot governed by 3 exclusion points
- Exponential tails (large & small exit times)
- Macroscopic observables ( $t$ ,  $S(t)$ ) reveals details of a topological constraint
- Knot relaxation governed by number of crossing points
- Athermal driving, yet, effective degrees of freedom randomized



# Outlook

- Different knot types
- Correlation between crossing points

**Many possibilities with granular chains**

# Entropic Tightening

with Matthew Hastings, Zahir Daya, Robert Ecke



- Equilibrium (counting states) prediction

$$P(n) \propto [n(N - n)]^{-d/2}$$

$$n/N \rightarrow 0 \quad \text{when} \quad N \rightarrow \infty$$

- Observed under nonequilibrium driving

**Role of entropy?**



Johnathan McKay

*My soul is an entangled knot  
Upon a liquid vortex wrought  
The secret of its untying  
In four-dimensional space is lying*

J. C. Maxwell