BOSTON UNIVERSITY

GRADUATE SCHOOL

Dissertation

KINETIC PROPERTIES OF STOCHASTIC PROCESSES

by

ELY BEN-NAIM

B.Sc., Hebrew University of Jerusalem, 1990

Submitted in partial fulfillment of the

requirements for the degree of

Doctor of Philosophy

1994

Approved by

First Reader

Sidney Redner, Ph.D.
Professor of Physics

Second Reader

Ganapathy Murthy, Ph.D.
Assistant Professor of Physics

to Ellen

ACKNOWLEDGMENTS

I would like to express my deepest gratitude to Sidney Redner, my thesis advisor. He has used his experience and knowledge to direct me to positive and creative avenues. He has engrained in me the lesson of seeking the beauty in science. I am also thankful for Sid's sincerity, patience and kindness.

I was fortunate to work with Paul Krapivsky over the past year, and I would like to thank him for all that he has taught me and for his friendship. I am grateful that I had the opportunity to collaborate with Daniel ben-Avraham, Francois Leyvraz, George Weiss and Jun Zhuo. I have enjoyed scientific discussions with Ronald Dickman, Alon Drory, Shlomo Havlin, Hernan Lerralda, Vladimir Privman, Stefan Schwartzer Haim Taitelbaum, Julian Talbot, Bill Young, Martin Vilela and Hwang Won Muk.

My stay in Boston would have never been the same without the friendship of Bill Hellman, Townsend Zwart, Ned Greene, Steve Harrington, Balaji Bashayam, Klaus Breuer and Monish Varma.

I would like to thank all my teachers, from elementary school through the Hebrew University. Special thanks to my High School Science teachers, Professor Levine and Professor Lipman for stimulating my interest in the Physical Sciences.

Last but not least, I thank my parents, Salomon and Simi, my siblings, Dina, Johnny and Alex and my neice, Lidi, for their love and for supporting me from afar.

I dedicate this thesis to Ellen, my wonderful wife, for being there for me and for her love.

KINETIC PROPERTIES OF STOCHASTIC PROCESSES

(Order No.)

ELY BEN-NAIM

Boston University Graduate School, 1994

Major Professor: Sidney Redner, Professor of Physics

ABSTRACT

In this thesis, kinetic and steady state properties of a range of stochastic and nonequilibrium models are investigated. A typical process is specified by a transport mechanism and a reaction mechanism and this leads naturally to simple "toy-like" models.

In the first part, the kinetics of ballistically-controlled reactions with continuous initial particle velocity distributions are studied. A non-universal power-law decay is found for both the concentration and the velocity, with the characteristic exponents governed by the form of the initial velocity distribution. For the case where particles annihilate upon contact, a mean-field theory provides good estimates for the decay kinetics. A ballistic aggregation model, inspired by one-lane traffic flows, is solved and the qualitative behavior of the reaction process is determined. The contribution of diffusion to the annihilation process with a bimodal initial velocity distribution is also studied. Using scaling arguments and series analysis techniques, it is shown that introduction of diffusion, even if small, alters the asymptotic behavior of the concentration of particles.

In the second part, a steady state model of the inhomogeneous two-species annihilation reaction, $A + B \to 0$, is presented. In this model, two different species A and B are injected with equal fluxes at the opposite boundaries of a finite system. Analytical results for the concentration profile are obtained by solving the reaction-diffusion equations. The dependence of the reaction rate on the flux is found and this result is applied to the time-dependent reaction model.

In the third part, collective properties of adsorption-desorption models in one dimension are studied. The steady state of a reversible parking process, where monodisperse particles adsorb and desorb on a one-dimensional substrate, is solved. The system exhibits a weak dependence on the adsorption rate in the desorption-controlled regime. To understand this behavior, an irreversible model incorporating immediate adsorption as well as desorption is introduced. A slow approach to the fully occupied state is found by heuristic arguments and by simulations. Furthermore, a cluster approximation technique is suggested to analyze the critical nature of a one-dimensional irreversible adsorption-desorption model.

Contents

List of Figures

List of Tables

Chapter 1

Introduction

"Would I find the complete many-body wave function, I would not know what to do with it" said Richard Feynman. This famous quote sums up my perspective of the essence of modern physics. To write a microscopic function that describes 10^{23} particles explicitly, a page larger than the sun-to-earth distance is required, thereby making such an approach of little practical use. Conceptual understanding of a physical system is necessary for making measurable predictions rather than obtaining a multivariative description of a many-particle system.

Disagreeing with Einstein's celebrated quote, I believe God's favorite toy is indeed the dice. Nature is often represented by stochastic terms. The quantum mechanical description of particles is essentially derived from probability amplitudes, which determine the likelihood of finding a particle at a certain point in time and space. Randomness governs not only quantum systems but also classical systems. The same mechanism that governs earnings in a dice throwing game controls the propagation of heat. As a result, heat propagates considerably slower in comparison with light.

Probabilistic descriptions are especially well-suited for analyzing many-particle phenomena. According to the central limit theorem, the statistical uncertainty diminishes as the number of random variables is increased. This uncertainty additionally diminishes when average macroscopic properties are calculated. Thus, it proves useful to study the limit of an infinite number of particles.

In this thesis, I study interacting particle systems by analyzing their underlying stochastic processes. A typical process is specified by a transport mechanism and an interaction mechanism, and this approach leads naturally to simple "toy-like" models. I focus on reaction processes and adsorption processes using mainly classical theoretical methods which are supported by computational studies. In addition, I present new theoretical methods and analysis techniques to investigate certain problems for which the conventional methods are not applicable.

In most of the problems we consider, time dependent phenomena are of interest, while in the remainder steady-state properties are important as well. In either situation, the asymptotic analysis of a properly defined microscopic process provides the desired physical insight. Generally, most details associated with the initial configuration vanish after waiting long enough or equivalently, become irrelevant as averaging procedures take place. Hence, we view the long time behavior as a tool for classifying processes into universality classes, *i. e.*, families that obey similar physics.

Generally, exact theoretical analysis is possible in the one-dimensional case only. However, one can project the understanding obtained for such soluble problems into a theory that holds in an arbitrary dimension. Scaling techniques are powerful in that respect, since they require a minimal amount of details and labor. In sufficiently high dimensions, our systems are expected to match a "mean-field" like behavior. To formulate such a theory we typically assume that all particles are neighbors of each other. This assumption yields a dimension-independent theory that ignores any spatial structures or microscopic fluctuations. The resulting deviation between the actual properties and the mean-field theory measures the extent to which particle interact. Specifically, because of effective repulsions between the particles our reactive systems exhibit anomalously slow kinetics. To a certain extent, a reacting particle system organizes itself such that the number of remaining reactants is maximized.

In equilibrium transitions between different states of the system obey the detailed balance condition that the flux between any two microscopic configurations of the system vanish. An equilibrium process also requires such transition rates to be microscopically reversible. Some of the adsorption processes we study are irreversible and hence, they are far from equilibrium. Not all possible configurations are accessible, resulting in significantly different steady state and kinetic properties than their equilibrium counterparts. In addition, collective behavior of the adsorbants cause slower than mean field approach to the steady state.

In chapter 2, a short review of some elementary kinetic processes is presented. The purpose of this mostly self-contained presentation is to place the work we have done in proper context and also to provide theoretical background. The starting point is a time-dependent solution of the Ising chain. We make use of the isomorphism between the spin system and annihilating random walkers to solve both processes. The correspondence between the slow decay kinetics of the diffusive single-species annihilation process and the Ising chain at its zero temperature critical point is established. Also, reversible adsorption is mapped to the equilibrium spin model at arbitrary temperature. Steady state properties are compared with the irreversible adsorption case, exemplifying the role of reversibility.

In chapters 3–5, the kinetics of reaction processes with ballistic particles are studied. Such a process can describe charged exciton recombination kinetics in an external electric field. The single species annihilation process, $A + A \to 0$, and the aggregation process, $A + A \to A$, with a continuous initial particle velocity distribution are studied. For the annihilation case, time-dependent properties of the concentration of particles and the typical velocity in one and two dimensions are investigated. A non-universal power-law decay is found for both the concentration and the velocity, in which the characteristic exponents, are governed by the form of the initial velocity distribution. Using scaling techniques and numerical integration of the Boltzmann equation, the exponents of the power law decay are estimated. Monte Carlo simulations of the process are performed and good agreement with the mean-field predictions is found.

The aggregation process, motivated by one-lane traffic flows, is useful in explaining the non-universal decays found for the annihilation model. Here aggregates move with a velocity equal to the minimal particle velocity, and thus, statistics of extremes govern this process. Asymptotically exact expressions are obtained for the velocity distribution and the joint mass-velocity distribution. Although the actual quantitative behavior of the ballistic aggregation and annihilation processes are different, both exhibit the same functional dependence on the initial conditions. Generalization of this model to situations with input and to spatially inhomogeneous situations are formally solved. For the input case, a transition to a steady state is also found as the input velocities are varied.

The contribution of diffusion to the annihilation process with a bimodal initial velocity distribution is also studied. Using scaling arguments, we show that introduction of diffusion, even if small, alters the asymptotic behavior of the concentration of particles. To confirm this prediction, a new general analysis method is developed. The approach is based on using an enumeration of all states to determine the exact time evolution of a reactive system. A Padé analysis technique is suggested for obtaining the asymptotic behavior of a process from its short-time Taylor series. This method is applied to ballistic annihilation problem with superimposed diffusion and the result of the scaling approach is verified.

In chapter 6, a steady state model of the inhomogeneous two-species annihilation reaction, $A + B \to 0$, is presented. This reaction scheme is relevant to studying electron-hole annihilation in irradiated semiconductors, for example. In our model, two different species A and B are injected with equal flux at the opposite boundaries of a finite system. The transport mechanism is diffusive and annihilation occurs when two different species meet. Analytical results for the concentration profile are obtained by solving the reaction-diffusion equations. The dependence of the reaction rate on the flux is found, and this result is applied to the time-dependent reaction model. The local reaction rate exhibits a transition from maximum near the boundary to maximum in the center of the system as the flux is increased. The solution technique is applied to systems with radial geometries in two- and three-dimensions as well.

In chapters 7 and 8, collective properties of adsorption-desorption models in one dimension are studied. A reversible parking process where monosize particles adsorb on a substrate is suggested for modeling binding of protein macromolecules on a surface. Exact results for the density and the distribution function of empty intervals in the steady-state are obtained. The system exhibits a weak dependence on the adsorption rate in the desorption-controlled regime. To understand this behavior, an irreversible model incorporating immediate adsorption as well as desorption is introduced. A slow approach to the fully occupied state is found by heuristic arguments, and this behavior is confirmed by Monte-Carlo simulations.

In chapter 8, an approximate solution to the contact process is developed. In this lattice model, particles desorb homogeneously and adsorb near occupied sites only. Such an autocatalytic reaction is closely related to surface oxidation of carbon monoxide on metal surfaces. This nonequilibrium process possesses an absorbing state and exhibits a continuous phase transition between an active state and an absorbing state. Such phase transitions are characterized by nontrivial critical exponents. In our approximation scheme, the correlation between two empty intervals is neglected. This approach yields different critical behavior, namely, a first order phase transition. However, for a significant range of desorption rates, the density and the gap distribution, that arises from our theory approximates the actual process accurately. Moreover, the values of the transition point and the critical exponents are close to the values found by series expansion techniques.

Chapter 2

Review

To put the work presented in this thesis in perspective, I present a short review of simple stochastic processes in this chapter. I begin with perhaps the simplest microscopic process modeling a chemical reaction, diffusive-driven single species annihilation. This process can be analyzed by considering the kinetic Ising model at zero temperature and identifying domain walls with particles. The classical solution, obtained by Glauber, is outlined as a starting point and used to derive the solution to the annihilation model. This correspondence establishes a deep relation between the spin model as a stochastic process and the collective properties of many interacting random walkers.

Moreover, the finite temperature Ising model can be mapped to a reversible adsorption model. The equilibrium Ising model solution yields, with a suitably defined temperature, a steady state solution to the adsorption process. Irreversible adsorption is then introduced and used to explain the significance of reversibility in kinetic processes.

Some of the most interesting phenomena in nonequilibrium physics are induced by fluctuations. In applications to chemical reactions, a striking example is the two species annihilation process. We describe the solution to a ballistically-driven reaction and stress the role of the initial density fluctuations in the asymptotic limit. This argument can be repeated for obtaining a scaling solution to the two species annihilation model. It is shown that slow kinetics take place as a result of the formation of single species domains.

2.1 The Kinetic Ising Model

One of simplest interacting particles system, the spin 1/2 model suggested by Ising [1], successfully describes the ferromagnetic phase transition [2]. In this model the interaction between nearest neighbors only is taken into account according to the following Hamiltonian,

$$H = -J \sum_i \sigma_i \sigma_{i+1}, \tag{2.1}$$

written explicitly for one dimension, with $\sigma_i = \pm 1$. When the spins are coupled to a heat reservoir, the equilibrium properties are unique, given the energy function of Eq. (2.1). The non-equilibrium behavior, however, depends on the dynamic leading the system to equilibrium and is generally not well defined. To define the time-dependent process, the transition rates between possible states of the system need to be specified. One can introduce sequential dynamics, *i. e.*, spins can change only one at a time. This yields a Markoff process, and taking these rates according to the detailed balance requirements of the equilibrium state allows a solution [3].

A many-spin system can be generally described by the collective probability function $P(\{\sigma\},t) \equiv P(\{\sigma_1,\ldots,\sigma_N\},t)$. When the individual spins, σ_i, undergo a spin-flip with rate $w_i(\sigma_i)$, the collective probability obeys the following rate equation,

$$\frac{\partial P(\{\sigma\},t)}{\partial t} = -\left(\sum_i w_i(\sigma_i)\right) P(\{\sigma\},t) + \sum_i w_i(-\sigma_i) P(\{\sigma_1,\cdots -\sigma_i,\cdots \sigma_N\},t). \tag{2.2}$$

The negative term represents loss of probability due to all possible transitions out of the state $\{\sigma\}$, while the gain term is obtained by noting that only configurations differing from $\{\sigma\}$ by one spin exactly contribute. One can check that summation of the rate equations over all possible states yields zero, since the total probability is conserved. We are interested in the evolution of average quantities such as the average spin value $\langle \sigma_i(t) \rangle = \sum_{\{\sigma\}} \sigma_i P(\{\sigma\},t)$ and the two-spin correlations $\langle r_{i,j}(t) \rangle =$

$\sum_{\{\sigma\}} \sigma_i \sigma_j P(\{\sigma\}, t)$. The rate equation for the single spin correlation is given by a summation of Eq. (2.2),

$$\frac{\partial\langle\sigma_i(t)\rangle}{\partial t} = \langle\sigma_i(t) w_i(\sigma_i(t))\rangle. \tag{2.3}$$

Similarly, the pair correlation function satisfies the rate equation

$$\frac{\partial\langle\sigma_i(t)\sigma_j(t)\rangle}{\partial t} = -2\langle\sigma_i(t)\sigma_j(t)\left[w_i(\sigma_i(t)) + w_j(\sigma_j(t))\right]\rangle. \tag{2.4}$$

To define the dynamics, the flip rates have to be specified. In analogy with the bilinear form of the Hamiltonian function, and under the assumption that the rates depend only on the values of the nearest neighbor spins, the following form was postulated by Glauber*

$$w_i(\sigma_i) = \frac{\alpha}{2}\left[1 - \frac{\gamma}{2}\sigma_i(\sigma_{i-1} + \sigma_{i+1})\right]. \tag{2.5}$$

These rates represent a tendency towards parallel spin alignment, for positive γ. In equilibrium, the probability that the system is in state $\{\sigma\}$ is given by the Boltzmann factor $\exp(-H(\{\sigma\})/k_B T)$, with k_B the Boltzmann constant and T the temperature. In addition, the equilibrium transition rates have to satisfy the detailed balance condition, namely, the flux between any two microscopic states of the system cancels. Therefore one finds

$$\frac{w_i(\sigma_i)}{w_i(-\sigma_i)} = \frac{p_i(-\sigma_i)}{p_i(\sigma_i)} = \exp\left(J\sigma_i(\sigma_{i-1} + \sigma_{i+1})/k_B T\right). \tag{2.6}$$

This relation is always fulfilled when the two neighboring spins are anti-parallel, while when they are parallel, the relation $(1+\gamma)/(1-\gamma) = \exp(4J/kT)$ is satisfied. After a simple manipulation, one finds

$$\gamma = \tanh(2J/k_B T). \tag{2.7}$$

Note that the prefactor α is arbitrary, since it enters through a mere scaling of the time variable. Indeed, a positive coupling constant J yields a positive γ, and therefore the spin system is derived towards a ferromagnetic state.

---

*The most general reflection-invariant bilinear form is actually $w_i(\sigma_i) = (\alpha/2)(1+\delta\sigma_{i-1}\sigma_{i+1})(1-(\gamma/2)\sigma_i(\sigma_{i-1} + \sigma_{i+1}))$, and Eq. (2.5) was chosen for simplicity.



Figure 2.1: The spin flip process

We are now in a position to write the explicit equations satisfied by the spin correlation function. The suggested form for the rates has the advantage that a closed set of equations emerges. We will concentrate on the pair correlations since it gives physical insight and understanding of both the kinetics and the non-equilibrium properties. We restrict ourself to the spatially homogeneous situation, where the spin correlation depends only on the distance between the two spins. Hence, we write $r_k(t) = \langle \sigma_i(t)\sigma_{i+k}(t)\rangle$. Using Eqs. (2.4) and (2.5), the rate equation is obtained as

$$\frac{\partial r_k(t)}{\partial(\alpha t)} = -2r_k(t) + \gamma\,(r_{k-1} + r_{k+1}) \qquad k \neq 0. \tag{2.8}$$

Curiously, the pair correlation satisfies a continuous-time discrete-space random walk differential equation with an annihilation rate $2(1-\gamma)$. Under the general initial conditions $r_k(0)$ and the boundary condition $r_0 = 1$, the pair correlation function takes the form, derived using generating functions in Appendix A,

$$r_k(t) = \eta^k + \sum_{l=1} \Big(r_l(0) - \eta^l\Big)\Big(I_{k-l}(2\alpha\gamma t) - I_{k+l}(2\alpha\gamma t)\Big), \tag{2.9}$$

with $\eta = (1 - \sqrt{1-\gamma^2})/\gamma = \tanh(J/kT)$.

The equilibrium solution $r_k^{eq} = \eta^k$ can be tacitly written in an exponential form,

$$r_k^{eq} = \exp(-k/\xi). \tag{2.10}$$

The constant ξ is readily identified as the correlation length, the spatial extent over which the spins are correlated. One can view this parameter as the typical size over which all spins are parallel. At the phase transition point, this length diverges. In the one-dimensional case, the critical temperature is zero, as seen from the relation $\xi = -1/\ln\ \tanh(J/kT)$. For finite temperatures, there is a finite correlation, indicating a paramagnetic behavior. In the limit of infinite temperature, the correlation length vanishes, and similarly the magnetization approaches zero.

The divergence of the correlation length is indicative of a critical slowing down mechanism, which can be seen by analyzing the temporal decay of the total magnetization. This quantity can be found by fixing one spin and summing over all correlations, (see Appendix A),

$$m(t) = m(0)\exp(-t/\tau) \qquad \text{with} \qquad \tau = \frac{1}{2\alpha(1-\gamma)} \tag{2.11}$$

As the system approaches the critical point, $\gamma \to 1$, the relaxation time τ diverges. This behavior is typical for systems at their transition point. Exponential temporal decay gives way to a slower power-law behavior. The nearest-neighbor correlation function, r_1, which is proportional to the average energy of the system, provides an example of such behavior. For simplicity, we consider the totally antiferromagnetic initial conditions, namely, $r_l(0) = (-1)^l$. Evaluating Eq. (2.9) under these initial conditions at zero temperature $\gamma = \eta = 1$ yields

$$r_1(t) = 1 - 2\exp(-2\alpha t) I_0(2\alpha t). \tag{2.12}$$

Since the modified Bessel functions have the asymptotic form $I_n(x) \sim e^x/\sqrt{2\pi x}$, the exponential factor cancels. One finds asymptotically, $1 - r_1(t) \sim (\pi\alpha t)^{-1/2}$. A power-law decay takes place at zero temperature, reminiscent of the general kinetic properties of systems at the critical point. This observation is significant to our studies, since we are mainly interested in kinetics of critical systems.

2.2 Diffusive Single-Species Annihilation

The model $A + A \to 0$ describes a chemical reaction in which diffusing particles annihilate upon contact [4, 5]. Diffusion is modeled by nearest-neighbor random walk, namely, particles hop with rate D to any nearest-neighbor site. An attempt to



Figure 2.2: Single species annihilation

land on an occupied site results in an annihilation event, as illustrated in Figure 2.2. The annihilation process occurs with a rate $2D$ since a hopping attempt of either particle in the correct direction enables a reaction. The kinetic Ising model at zero temperature can be mapped into this stochastic process by identifying antiparallel spins with particles and conversely, parallel ones with empty sites. For example, the spin configuration $\cdots\downarrow\uparrow\uparrow\downarrow\cdots$ corresponds to the particle configuration $\cdots AOA\cdots$. The annihilation process $AA \to OO$ corresponds to the spin flip process $\uparrow\downarrow\uparrow\to\uparrow\uparrow\uparrow$, and the forbidden creation process $OO \to AA$ corresponds to $\uparrow\uparrow\uparrow\to\uparrow\downarrow\uparrow$. To establish the isomorphism between the particle and the spin processes, the corresponding rates are matched. With the help of Figures 2.1 and 2.2, one has $\alpha(1+\gamma)/2 = 2D$ and $\alpha(1-\gamma)/2 = 0$, or equivalently, $\alpha = 2D$ and $\gamma = 1$. Note also that the hop process rate D matches the rate $\alpha/2$, which describes spin-flips that do not change the energy. At zero temperature, energy raising transitions are forbidden ($\gamma = 1$) or equivalently, no creation of particles are allowed in the particle process. The particle concentration, $c(t)$, equals the domain boundary concentration, and from the definition of the nearest neighbor correlation, one finds

$$c(t) = \frac{1 - r_1(t)}{2}. \tag{2.13}$$

Thus, the antiferromagnetic initial condition discussed in the previous section corresponds to the completely filled initial state. Finally, the time dependent concentration is found by substituting the proper rates in Eq. (2.11) to yield

$$c(t) = I_0(4Dt)\exp(-4Dt). \tag{2.14}$$

Interestingly, this concentration equals the probability that the displacement of a single random walk, hopping with rate $2D$, vanishes at time t. This can be partially understood by noting that the distance between two identical random walks itself executes a random walk with a hopping rate $2D$. In the long time limit, $t \gg 1/D$, the concentration shows a power law decay,

$$c(t) \sim (8\pi Dt)^{-1/2}. \tag{2.15}$$

Going back to $T = 0$ kinetic Ising model, the domain boundaries can be viewed as performing a random walk. When two domains meet, an annihilation event occurs and a combined larger domain form. The annihilating random walk argument sheds light on the behavior of the spin system.

We present now a heuristic argument that reproduces the above asymptotic result and which can also be easily generalized to higher dimensions. Let us assume that at given time, there exist an "interaction length", l, such that particles initially separated by a distance smaller than l have had a chance to interact (annihilate). Space can be divided now into "blocks" of volume l^d, and under this interaction length assumption, the number of surviving particles in each such block can be taken as one. The concentration is, thus, given by $c \sim l^{-d}$. Since the transport mechanism is diffusive, it is reasonable to assume that the interaction length is given by the average spread of a diffusive particle, $l \sim \sqrt{Dt}$. By substituting the time dependent length into the expression describing the concentrations we obtain

$$c(t) \sim (Dt)^{-d/2}. \tag{2.16}$$

Note that the concentration does not depend on the initial concentration. Consequently, the asymptotic form given in Eq. (2.15) is expected to hold for arbitrary random initial conditions.

Let us consider now the case where particles hop to a randomly chosen site, and annihilation occurs whenever an attempt to land on an occupied site is made. In this case, we ignore the spatial dimensionality, and we imagine that all sites are connected to each other. Thus, the concentration obeys the mean-field evolution equation $dc(t)/dt = -kc^2$, where k is the reaction rate, and whose solution is

$$c(t) = \frac{1}{c_0^{-1} + kt}. \tag{2.17}$$

At two dimensions, the block argument matches the mean-field behavior, and we conclude that critical dimension for this process is $d_c = 2$. Above this critical dimension, t^{-1} decay occurs. Furthermore, the slower than mean-field decay indicates an effective repulsion between particles. This repulsion is manifested by a vanishing interparticle distance distribution function [6]. This system is one example that illustrates the crucial role that spatial fluctuations play in low dimensions.

The block argument is a robust one; it remains the same for a class of microscopic interaction mechanism. For example, consider the coalescence process $A + A \to A$ [6], where the asymptotic form $(Dt)^{-1/2}$ should hold up to a prefactor. Indeed, the diffusion-limited fusion of photogenerated excitons in a one-dimensional tetramethylammonium manganese trichloride system was observed to obey this asymptotic decay [7].

As mentioned previously, the annihilating particles system is dual to an adsorption process. The mapping is simple; particles are identified with empty sites ("holes") and vice versa. Since a hop of a particle to the right, $AO \to OA$, is equivalent to a hop of a hole to the left, particle diffusion corresponds to hole diffusion. The annihilation process, on the other hand, $AA \to OO$, corresponds to adsorption of dimers. Hence, by reversing the time arrow, the annihilation process is mapped into the adsorption process. Therefore, the annihilation process is isomorphic to an adsorption process where pairs of particles adsorb on the surface with rate $2D$ and hop to nearest neighbor sites with rate D. In the adsorption process, the concentration, given by $1 - c(t)$, approaches unity with a power-law correction $t^{-1/2}$ [8]. In chapter 7, we discuss a closely related process occurring on the continuum where a logarithmic approach to the completely filled state occurs. It is also seen that irreversible adsorption, where particles are allowed to diffuse, is also isomorphic to the



Figure 2.3: The reversible adsorption process

kinetic Ising model. In the next section, we expand this approach to one-dimensional adsorption-desorption processes.

2.3 Adsorption Processes: Role of Reversibility

In a reversible adsorption process, particles adsorb on a surface and similarly, particles are also allowed to desorb. Typically, the adsorbing particles are r-mers, *i. e.*, they occupy r consecutive lattice sites. In this section, we discuss an elementary dimer adsorption-desorption process. For an adsorption attempt to be successful, both sites must be vacant. The aforementioned mapping to the kinetic Ising model is used here to obtain the equilibrium solution and the asymptotic approach towards this state [9]. In the reversible adsorption case, dimers can adsorb only if both lattice site are vacant. The two adsorbing particles then diffuse independently on the surface, as shown in Figure 2.3. The three rates defining this process are the adsorption rate k_+, the desorption rate k_-, and the hopping rate D. Using the isomorphism to the spin model, established in the previous chapter (see figure 2.1), we match the above rates with the spin-flip rates $\alpha(1+\gamma)/2$, $\alpha(1-\gamma)/2$ and $\alpha/2$, respectively. Consequently, the temperature of the equivalent spin model is given by $\exp(-4J/k_B T) = k$, with $k = k_+/k_-$. Note that the kinetic Ising model has only two independent parameters and hence, the hopping rate cannot be arbitrary and is in fact given by $D = k_+ + k_-$. Using the pair correlation of Eq. (2.10) and Eq. (2.11), the equilibrium concentration is found to be

$$c^{eq}(k) = \frac{\sqrt{k}}{1+\sqrt{k}}. \tag{2.18}$$

For small adsorption rates, the concentration is an algebraic function of the normalized adsorption rate $c \sim \sqrt{k}$. In contrast, for reversible adsorption of monomers, the solution is a trivial one and the corresponding dependence is a linear one. The temporal approach to the equilibrium solution is related to the decay of the energy fluctuations in the Ising model, since r_1 is proportional to the average energy. The relaxation time τ is identical to the one previously found for the relaxation of the magnetization $\tau = 1/2(1-\gamma)$, and as a result,

$$\tau = \frac{1+k}{4k}. \tag{2.19}$$

The relaxation time diverges as k vanishes, reminiscent of a critical point.

However, the qualitative nature of this behavior changes drastically when the transition rules are changed. For example, when adsorption of monomers only is coupled with desorption of dimers, the following results were found, $c \sim k^{1/3}$ and $\tau \sim k^{-2/3}$ for $k \ll 1$. This result is obtained by noting that the monomer adsorption process corresponds to flipping all spins to the right of the adsorbing sites [9]. The two process are fundamentally different. While the former is reversible, the latter is not. Therefore, the steady state obtained in the monomer adsorption case is not an equilibrium one (detailed balance is not satisfied). Typically, most nonequilibrium process lead to a steady state that does not satisfy the detailed balance condition. This example illustrates that reversibility is a fundamental aspect of a stochastic process.

The desorption limited case sheds light on the importance of reversibility. While the concentration in the reversible case approaches unity for $k \to \infty$, non-trivial behavior occurs when adsorption only occurs, namely, irreversible adsorption. In this problem dimers are deposited randomly on an initially empty surface. An adsorption event is successful only if both sites are empty. With the empty initial conditions, the system reaches a jammed configuration, where no further adsorption events are possible. The steady-state density is given by the famous Flory result [10]

$$c_\infty = 1 - e^{-2}. \tag{2.20}$$

In appendix B, we detail a solution for the general for k-mer random sequential adsorption process in one-dimension, and Eq. (2.20) follows from the $k = 2$ solution. Starting from the partially occupied state with the dimers separated by a single vacant site, the initial state is left unchanged, and the density is always 2/3. This discrepancy shows that in the irreversible adsorption case the limiting concentration depends strongly on the initial conditions.

2.4 Fluctuation Dominated Reactions

In this section we solve for the concentration in a ballistic annihilation process [11]. The solution is then used to explain the significance of initial spatial fluctuations in determining the asymptotic kinetics. The similarity to diffusively-driven two-species annihilation is stressed using a heuristic argument.

The single-species ballistic annihilation process is defined as follows. Ballistic particles are distributed randomly with density c_0 on the continuum. Particles move with constant velocity, and a collision results in the annihilation of both colliding particles. Here we discuss the case where particles are assigned two opposite velocities: $+v_0$ with probability p_+, and similarly, $-v_0$ with probability $p_- = 1 - p_+$.

In the two-velocity case it is possible to identify all collision partners, given the initial positions of the particles. The collision times are then determined by incorporating the distances between the particles. Two particles can annihilate only if they are separated by an even number of particles. Denoting particles by their order, $k = 1, \ldots, 2n$, we define b_n as the probability that all such $2n$ particles eventually react among themselves, with $b_0 = 1$. For example, $b_1 = p_+ p_-$. Taking into account all possible collision partners of the $k = 1$ particle gives a recursion relation for $n > 1$,

$$b_n = p_+ p_- \sum_{k=0}^{n-1} b_k b_{n-1-k}. \tag{2.21}$$

By introducing the generating function $B(z) = \sum_{n=0} b_n z^n$ and summing Eq. (2.21) over all n, one finds

$$B(z) = \frac{1 - \sqrt{1 - 4p_+p_-z}}{2p_+p_-z}, \tag{2.22}$$

and by expanding $B(z)$ in a power-series one has

$$b_n = \frac{(p_+p_-)^n(2n)!}{n!(n+1)!}. \tag{2.23}$$

Starting with an unequal concentration of right moving and left moving particles, the system reaches a state with the majority species only present, $c_\infty = c_0|p_+ - p_-|$. We now obtain the concentration for the uniform initial condition, where the separation between two neighboring particles is always $1/c_0$. To survive to time t, a particle has to avoid collisions with all k-particles such that $|k| > 2c_0v_0t$. Using the reaction probabilities, the time dependent concentration is written as $c(t) - c_\infty = 2c_0p_+p_- \sum_{n=[c_0v_0t]} b_n$. Substituting the actual reaction probability given by Eq. (2.23) in this expression yields the long time behavior of the concentration. When the initial concentration of the species are not equal, the concentration approaches the final state as $c(t) - c_\infty \sim (c_0v_0t)^{-3/2}\exp(-\text{const.}\cdot c_0v_0t)$. In contrast, when $p_- = p_+ = 1/2$, a power-law decay is found,

$$c(t) \sim \left(\frac{c_0}{2\pi v_0 t}\right)^{1/2}. \tag{2.24}$$

Despite the apparent similarity of the result with that of the diffusively-driven single-species annihilation, the underlying mechanism for the decay is different. While the corresponding $(Dt)^{-1/2}$ decay is independent of the initial conditions, the concentration of Eq. (2.24) exhibits a dependence on c_0.

A heuristic argument, similar to the one used describing the $A + A \to 0$ process, can be applied to determine the decay of the density in the ballistic case. We assume that within an interaction length l all $N = N_+ + N_-$ initially present particles have had a chance to interact. The total number of surviving particles will thus be $\Delta N = |N_+ - N_-|$. Using the central limit theorem, $\Delta N \sim \sqrt{N}$, and consequently, the asymptotic concentration is found

$$c \sim \frac{\sqrt{N}}{l} \sim \left(\frac{c_0}{l}\right)^{1/2}. \tag{2.25}$$

Substituting the interaction length $l \sim v_0 t$, the exact result of Eq. (2.24) is reproduced. It is seen that the system arranges into domains of particles moving with identical velocity. This spatial organization enhances the survival probability by a shielding mechanism. A right moving particle is protected from annihilation by all consecutive right moving particles placed to its right. Inspired in part by these initial spatial fluctuations effects, we consider in chapter 3 ballistic annihilation with continuous initial velocity distributions.

Another important process that exhibits strong dependence on initial fluctuations is the diffusively-driven two-species annihilation [12, 13, 14]. Here two different species, denoted by A and B, are randomly distributed initially with equal concentrations. Particles move diffusively, and reactions are allowed only between opposite species. The fluctuation argument, presented for the ballistic reaction case, is applicable here, and a generalization of Eq. (2.25) gives $c \sim (c_0/l)^{d/2}$. The interaction length grows according to $l \sim \sqrt{Dt}$ since the transport mechanism is diffusive. Consequently, the following asymptotic form of the concentration is found

$$c(t) \sim \left(\frac{c_0^2}{Dt}\right)^{d/4}. \tag{2.26}$$

The concentration crosses over to the mean-field behavior at $d_c = 4$. Again, the slower than mean-field behavior is made possible by formation of ever growing domains occupied by one species only.

All the soluble processes we presented in this chapter are deeply related to the theory of random walks. For example, b_n, the probability describing the ballistic annihilation process is identical to the probability that a biased random walker returns to its starting point for the first time at the n^{th} step. Furthermore, the concentration in the diffusive driven annihilation equals the probability that a continuous time random walk is at the origin at time t. Curiously, however, the transport mechanism in the ballistic reaction process is not a diffusive one.

Chapter 3

Non-Universal Decay Kinetics in Ballistic Annihilation

A number of interesting results have been recently reported for the kinetics of irreversible agglomeration. In an attempt to describe phenomena such as fluid vortices coalescence [15] and planet formation by accretion [16], the following idealized aggregation model has been introduced. In this process, written symbolically as $A_i + A_j \to A_{i+j}$, the trajectories are ballistic, and the collisions between aggregates conserve momentum [17, 18]. The subscripts i, j refer to the (conserved) mass of the aggregates. A scaling argument suggests that the concentration of aggregates decays as $t^{-\alpha}$, with $\alpha = 2d/(d+2)$, where d is the spatial dimension [17]. This dimension dependence for all d is atypical of the behavior pattern exhibited by diffusion-controlled reactions, as discussed in the review chapter. Moreover, we have shown in section 2.4 that the ballistic annihilation process $A + A \to 0$ with two opposite particle velocities exhibits anomalously slow decay kinetics. This behavior was accounted for by the formation of large-scale spatial heterogeneities, a mechanism similar to the one governing the diffusive-driven two-species annihilation [12, 14, 19, 20, 21]. This spatial organization enhances the survival probability of a single particle.

Motivated in part by these intriguing features, we investigate the decay kinetics of the elementary single-species annihilation process, $A + A \to 0$, for arbitrary continuous initial velocity distributions. Such processes describe actual nonequilibrium processes such as electron-hole recombination and the decay of excitations induced by irradiation [22, 23]. We find that the decay of the density strongly depends on the form of the initial velocity distribution. Moreover, faster particles annihilate at a higher rate, leading to a decay of the typical velocity as well.

We suggest a scaling argument, based on the typical collision time, that relates the concentration to the typical velocity. A Boltzmann equation for the evolution of the velocity distribution verifies this relation. Using the anticipated scaling forms of both the velocity and the velocity distribution, we eliminate the time variable from this equation. We combine the limiting behaviors of the velocity distribution into an ansatz solution. This solution suggests nonuniversal dependence of α, the exponent describing the concentration decay, on the details of the initial velocity distribution near the origin. In one and two dimensions, numerical integration of the Boltzmann equation and Monte Carlo simulations verify our predictions. The Boltzmann equation yields values of the decay exponents that are in good agreement with the actual simulations. We conclude that the success of this mean-field approximation is made possible by the relatively small velocity correlations between neighboring particles.

3.1 Scaling Properties

At time $t = 0$, the system consists of identical particles which are distributed in space with $P(v, t = 0)$ the initial concentration of particles of velocity v. For simplicity, only symmetric distributions with respect to the origin are studied. The decay kinetics appear to be independent of the initial spatial distribution of particles, and thus we focus on a random initial distribution of interparticle distances. Particles move according to their initial velocity until a collision occurs, which results in the removal of both colliding particles, as illustrated in Figure 3.1. We are interested in determining the time dependence of the macroscopic concentration, $c(t) = \int dv P(v, t)$, and the moments of the velocity distribution, $\langle v^n \rangle^{1/n} = (\int dv\, v^n P(v, t)/c(t))^{1/n}$.

Figure 3.1: Illustration of the annihilation process

A simple power counting argument relates the exponents which characterize the decay of the concentration with the exponent corresponding to the typical velocity. Consider a system of identical particles of fixed radius r at concentration c which move with a velocity of the order of $v_{\rm rms}$. From an elementary mean-free path argument, the time between collisions is $t \sim 1/cv_{\rm rms}r^{d-1}$. If one assumes the following power law forms for the concentration and $v_{\rm rms}$,

$$c \sim t^{-\alpha} \qquad v_{\rm rms} \sim t^{-\beta}, \tag{3.1}$$

then the mean-free path argument indicates that the relation $\alpha + \beta = 1$ should hold for all spatial dimension d.

Since the lifetime of particles with velocity v is proportional to $1/v$, faster particles tend to annihilate more quickly, and the typical velocity should decay in time. By the relation between α and β, a value of α less than unity is therefore implied. We further argue that there is a strong dependence of the exponent α on the form of the initial velocity distribution. This behavior differs from the mean field prediction of $c \sim t^{-1}$ which arises from the naive rate equation $\dot{c} \propto -kc^2$.

3.2 The Boltzmann Equation

A useful approach for determining the decay kinetics is to write a Boltzmann equation for the time evolution of the velocity distribution. For simplicity, consider the case of one dimension; generalization to higher dimensions follows naturally. Let $P(x, v, t)$ be the density of particles with velocity v at position x and at time t. At time $t + \Delta t$, the velocity distribution changes both because of translation of particles and because of reactions. We treat the reaction term in a mean-field approximation by assuming that a particle at $x' < x$ and velocity $v' > v$ will necessarily react with the target particle at (x, v) when $x - x' < (v' - v)\Delta t$. There is a complementary contribution due to collisions between the target and a particle located at $x' > x$ with $v' < v$.

The sum of these two contributions leads to the Boltzmann equation for ballistic annihilation

$$\begin{aligned} &P(x + v\Delta t, v, t + \Delta t) - P(x, v, t) = \\ &-kP(x, v, t)\Big[\int_{v}^{\infty} \int_{x-(v'-v)\Delta t}^{x} dx'\, P(x', v', t) + \int_{-\infty}^{v} dv' \int_{x}^{x+(v-v')\Delta t} dx'\, P(x', v', t)\Big], \end{aligned} \tag{3.2}$$

where k is a dimensionless reaction constant. Since a collision leads to particle annihilation, there is no collision-induced gain term in the equation. This approximate equation overcounts collisions, since the incident particle at x' may react with a third particle rather than with the target particle.

To analyze the Boltzmann equation, we expand Eq. (3.2) to first order in Δt to arrive at

$$\frac{\partial P(x, v, t)}{\partial t} = -v\frac{\partial P(x, v, t)}{\partial x} - kP(x, v, t)\int_{-\infty}^{\infty} dv'|v - v'|P(x, v', t)\,. \tag{3.3}$$

Since the initial velocity distribution is spatially homogeneous, and because of the homogeneity of the reaction process, we assume that the velocity distribution remains spatially homogeneous. Thus we write $P(v, t)$ to signify the time-dependent and spatially homogeneous density of particles with velocity v. This implies that the convective term $\partial P/\partial x$ in the Boltzmann equation may be set to zero, leading to

$$\frac{\partial P(v, t)}{\partial t} = -kP(v, t)\int_{-\infty}^{\infty} dv'\,|v - v'|P(v', t)\,. \tag{3.4}$$

Thus the $|v - v'|$ dependence of the integral kernel controls the reaction rate between two particles. Eq. (3.4) is also strongly reminiscent of the Smoluchowski equation for ballistic aggregation [18]. Despite the uncontrolled nature of the approximations underlying Eq. (3.4), this formulation gives a useful quantitative description of the decay kinetics.

The first step in our analysis of the Boltzmann equation is to apply dimensional analysis, together with the assumed asymptotic behaviors, $c \sim t^{-\alpha}$ and $v_{\rm rms} \sim t^{-\beta}$, to reduce Eq. (3.4) to a single variable equation. From these considerations, we expect that the velocity distribution will have the following scaled form

$$P(v,t) = \frac{c_0}{v_0}\left(\frac{t}{t_0}\right)^{\beta-\alpha} f(z), \tag{3.5}$$

where the scaling function $f(z)$ depends only on the dimensionless velocity

$$z = \frac{v}{v_0}\left(\frac{t}{t_0}\right)^{\beta}. \tag{3.6}$$

Here $t_0 = 1/(kc_0v_0)$ is the initial characteristic time between reactions, with k the dimensionless reaction constant, c_0 the initial concentration, and v_0 the initial rms velocity.

Upon substituting this scaling form into Eq. (3.4), we confirm the anticipated exponent relation by equating all the resulting temporal prefactors

$$\alpha + \beta = 1. \tag{3.7}$$

Additionally, we obtain a rescaled Boltzmann equation for the scaling function

$$(2\beta - 1)f(z) + \beta z f'(z) = -f(z)\int_{-\infty}^{\infty} dz'\,|z - z'|\,f(z'). \tag{3.8}$$

Notice that this equation is invariant under the transformation $f(z) \to a^2 f(az)$, so that a unit normalization of $f(z)$ can be achieved by a scale change in z. We observe that the scaling form $f(z)$ depends heavily on the value of β, a behavior that was confirmed by direct numerical integration of the Boltzmann equation.

To find the tail of the scaled velocity distribution, we approximate $|z - z'| \sim |z|$ in the integral and use the fact that $f(z)$ vanishes as $z \to \infty$. These steps reduce Eq. (3.1) to a simple differential equation whose solution is

$$f(z) \sim |z|^{(1-2\beta)/\beta} \exp(-|z|/\beta), \qquad |z| \gg 1. \tag{3.9}$$

Near the origin, Eq. (3.8) admits different solutions for $f(z)$, and correspondingly, the exponent β depends on the form of the initial velocity distribution.

To treat the small-z limit, we first divide Eq. (3.8) by $f(z)$ to yield

$$(2\beta - 1) + \beta \lim_{z\to 0} z\,(\ln f(z))' = -2\int_0^\infty dz'\,|z'|\,f(z'). \tag{3.10}$$

Consider an initial velocity distribution with a power-law tail in the small velocity limit, $P(v,t=0) \propto |v|^\mu$. Our Monte Carlo simulations (discussed below) reveal that the scaled distribution retains the same power-law form, $f(z) \sim z^\mu$, in the small-z limit. By adopting this form in Eq. (3.10), the second term is simply equal to $\beta\mu$. The resulting equation then predicts that β is a monotonically decreasing function of μ (for $\mu > -1$) whose precise form depends on the first moment of $f(z)$. This moment, in turn, depends on the full details of the velocity distribution. For example, if we take as a trial function $f(z) \propto z^\mu e^{-z/\beta}$ in Eq. (3.10), *i. e.*, the product of the asymptotic behaviors, we obtain $\beta = 1/(3+2\mu)$. This estimated form qualitatively mirrors values for β obtained by numerical integration of Eq. (3.4) and on Monte Carlo simulations for various values of μ.

The generalization of the scaling approach for the Boltzmann equation to higher spatial dimensions is straightforward. The scaled velocity distribution function now takes the form

$$P(v,t) = \frac{c_0}{v_0}\left(\frac{t}{t_0}\right)^{\beta d-\alpha} f(\vec{z}\,), \quad \text{with} \quad \vec{z} = \frac{\vec{v}}{v_0}\left(\frac{t}{t_0}\right)^{\beta}, \tag{3.11}$$

where the exponent combination βd originates from an integration over d-dimensional velocity space. The corresponding rescaled equation for $f(\vec{z}\,)$ becomes

$$((d+1)\beta - 1)f(\vec{z}\,) + \beta\vec{z}\cdot\nabla f(\vec{z}\,) = -f(\vec{z}\,)\int_{-\infty}^{\infty} d\vec{z}\,'|\vec{z} - \vec{z}\,'|f(\vec{z}\,'). \tag{3.12}$$

We emphasize that for higher dimensions, the Boltzmann equation involves a qualitatively different type of approximation in comparison with the one-dimensional case. To determine the relation between the exponents μ and β, we again assume a small-z power law form $f(\vec{z}\,) \propto |\vec{z}\,|^\mu$, with $\mu > -d$, and a relatively sharp cutoff in $f(\vec{z})$ for large $|\vec{z}\,|$. When employed in the Boltzmann equation, these assumptions again lead to a qualitatively correct μ-dependence of β. Namely, β monotonically decreases with μ, and has the limits $\beta \to 1$ for $\mu \to -d$ and $\beta \to 0$ for $\mu \to \infty$. As in the case of one dimension, if we adopt $f(\vec{z}\,) \propto |\vec{z}\,|^{\mu} e^{-|\vec{z}\,|/\beta}$, then Eq. (3.12) gives the $\beta = 1/(1 + 2d + 2\mu)$. Conversely, as the spatial dimension increases, β decreases systematically, and the limit $\alpha = 1$ is approached but never reached. This behavior corresponds to the naive rate equation $\dot{c} = -kc^2$. Thus only in the $d = \infty$ limit are particle trajectories sufficiently transparent that the typical velocity does not decrease. This is in contrast to the situation of many diffusion-controlled reactions for which "transparent" behavior occurs for only $d >= d_c$ with d_c finite [14].

3.3 Numerical Analysis

To test our approximate analysis, we have performed direct numerical integration of the Boltzmann equations, Eq. (3.4) and Eq. (3.12) (see Table 3.1). In one dimension, a typical integration was based on dividing the velocity range $[-1, 1]$ into 200 bins with a time step of $\Delta t = 0.15$. A finer level of resolution gives essentially identical results. The integration was performed to 1000 time steps. To estimate the value of β, we computed the "test" scaling function $f(z;t)_{\rm test} \propto t^{1-2\beta_{\rm test}} P(v,t)$ at different times and adjusted $\beta_{\rm test}$ to achieve the best data collapse between different data sets. By this method, we obtained estimates for the exponent β with an uncertainty of less than 0.005.

We also performed Monte Carlo simulations for ballistic annihilation in one and two dimensions which are based on two independent approaches. One method simulates the ballistic motion as a biased random walk. In two dimensions, a particle at $\vec{r}_i$ is assigned a velocity (v_{ix}, v_{ix}), with $|v_{ix}|, |v_{iy}| < 1$, according to an initial velocity distribution with support in $[-1, 1]^2$. A move attempt consists of picking each occupied site $\vec{r}_i$ at random, and the particle is moved by an amount $(\mathrm{sgn}(v_{ix}), 0)$ with probability $|v_{ix}|$ and by an amount $(0, \mathrm{sgn}(v_{iy}))$ with probability $|v_{iy}|$. If a particle lands on an occupied site, both particles are removed from the system. The time is then incremented by the inverse of the current number of particles in the system. The attraction of this stochastic method is that it is easily implemented