

Nontrivial Exponents in Records Statistics

Eli Ben-Naim

Los Alamos National Laboratory

with: Paul Krapivsky (Boston), Pearson Miller (Yale)

E. Ben-Naim and P.L. Krapivsky, arXiv:1305:4227

P. W. Miller and E. Ben-Naim, arXiv: 1308:xxxxx

Talk, paper available from: <http://cnls.lanl.gov/~ebn>

Deep Computation in Statistical Physics, Santa Fe Institute, August 9, 2013

Plan


- I. Incremental records
- II. Superior records
- III. Inferior records
- IV. General distribution functions
- V. Earthquake data

Motivation

- Weather: record high & low temperatures Havlin 03
- Finance: stock prices Bouchaud 03
- Insurance: extreme/catastrophic events Embrechts 97
- Evolution: growth rate of species Krug 05
- Sports
- First passage phenomena Redner 01
Majumdar 13

First passage properties of extreme statistics

Incremental Records



Incremental sequence of records

every record improves upon previous record by yet smaller amount


random variable = $\{0.4, 0.4, 0.6, 0.7, 0.5, 0.1\}$

latest record = $\{0.4, 0.4, 0.6, 0.7, 0.7, 0.7\}$ \uparrow

latest increment = $\{0.4, 0.4, 0.2, 0.1, 0.1, 0.1\}$ \downarrow

What is the probability all records are incremental?


Probability all records are incremental



$$S_N \sim N^{-\nu} \quad \nu = 0.31762101$$

Power law decay with nontrivial exponent
Question is free of parameters!

Uniform distribution



- The variable x is randomly distributed in $[0:1]$

$$\rho(x) = 1 \quad \text{for} \quad 0 \leq x \leq 1$$

- Probability record is smaller than x

$$R_N(x) = x^N$$

- Average record

$$A_N = \frac{N}{N+1} \quad \implies \quad 1 - A_N \simeq N^{-1}$$

- Number of records

$$M_N = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{N}$$

Distribution of records

- Probability a sequence is inferior and record $< x$

$$G_N(x) \implies S_N = G_N(1)$$

$$x_2 = x_1$$


- One variable

$$G_1(x) = x \implies S_1 = 1$$

$$x_2 = 2x_1$$

- Two variables

$$x_2 - x_1 > x_1 \quad G_2(x) = \frac{3}{4} x^2 \implies S_2 = \frac{3}{4}$$



- In general, conditions are scale invariant $x \rightarrow ax$
- Distribution of records for incremental sequences


$$G_N(x) = S_N x^N$$

- Distribution of records for all sequences equals x^N

Statistics of records are standard

Fisher-Tippett 28
Gumbel 35

Scaling behavior



- Distribution of records for incremental sequences

$$G_N(x)/S_N = x^N = [1 - (1 - x)]^N \rightarrow e^{-s}$$

- Scaling variable

$$s = (1 - x)N$$

Exponential scaling function

Distribution of increment+records

- Probability density $S_N(x,y)dx dy$ that:
 1. Sequence is incremental
 2. Current record is in range $(x,x+dx)$
 3. Latest increment is in range $(y,y+dy)$ with $0 < y < x$

- Gives the probability a sequence is incremental

$$S_N = \int_0^1 dx \int_0^x dy S_N(x, y)$$

- Recursion equation incorporates memory

$$S_{N+1}(x, y) = \underbrace{x S_N(x, y)}_{\text{old record holds}} + \int_y^{x-y} dy' \underbrace{S_N(x - y, y')}_{\text{a new record is set}}$$

- Evolution equation includes integral, has memory

$$\frac{\partial S_N(x, y)}{\partial N} = -(1 - x) S_N(x, y) + \int_y^{x-y} dy' S_N(x - y, y')$$

Scaling transformation

- Assume record and increment scale similarly

$$y \sim 1 - x \sim N^{-1}$$

- Introduce a scaling variable for the increment

$$s = (1 - x)N \quad \text{and} \quad z = yN$$

- Seek a scaling solution

$$S_N(x, y) = N^2 S_N \Psi(s, z)$$

- Eliminate time out of the master equation

$$\left(2 - \nu + s + s \frac{\partial}{\partial s} + z \frac{\partial}{\partial z} \right) \Psi(s, z) = \int_z^\infty dz' \Psi(s + z, z')$$

Reduce problem from three variables to two

Factorizing solution

- Assume record and increment decouple

$$\Psi(s, z) = e^{-s} f(z)$$

- Substitute into equation for similarity solution

$$\left(2 - \nu + s + s \frac{\partial}{\partial s} + z \frac{\partial}{\partial z}\right) \Psi(s, z) = \int_z^\infty dz' \Psi(s + z, z')$$

- First order integro-differential equation

$$z f'(z) + (2 - \nu) f(z) = e^{-z} \int_z^\infty f(z') dz'$$

- Cumulative distribution of scaled increment $g(z) = \int_z^\infty f(z') dz'$

- Convert into a second order differential equation

$$z g''(z) + (2 - \nu) g'(z) + e^{-z} g(z) = 0 \quad \begin{array}{l} g(0) = 1 \\ g'(0) = -1/(2 - \nu) \end{array}$$

Reduce problem from two variable to one

Distribution of increment

- Assume record and increment decouple

$$zg''(z) + (2 - \nu)g'(z) + e^{-z}g(z) = 0 \quad \begin{array}{l} g(0) = 1 \\ g'(0) = -1/(2 - \nu) \end{array}$$

- Two independent solutions

$$g(z) = z^{\nu-1} \quad \text{and} \quad g(z) = \text{const.} \quad \text{as} \quad z \rightarrow \infty$$

- The exponent is determined by the tail behavior

$$\nu = 0.31762101$$


- The distribution of increment has a broad tail

$$P_N(y) \sim N^{-1}y^{\nu-2}$$

Increments can be relatively large
problem reduced to second order ODE

Numerical confirmation

Monte Carlo simulation versus integration of ODE




Increment and record become uncorrelated

Superior Records

- Start with sequence of random variables

$$\{x_1, x_2, x_3, \dots, x_N\}$$



- Calculate the sequence of records

$$\{X_1, X_2, X_3, \dots, X_N\} \quad \text{where} \quad X_n = \max(x_1, x_2, \dots, x_n)$$

- Compare with the expected average

$$\{A_1, A_2, A_3, \dots, A_N\} = \{1/2, 2/3, 3/4, \dots, N/(N+1)\}$$


- Superior sequence = records always exceeds average

$$X_n > A_n \quad \text{for all} \quad 1 \leq n \leq N$$

- What fraction S_N of sequences is superior?

measure of “performance”

Numerical simulations



$$S_N \sim N^{-\beta}$$

$$\beta = 0.4503 \pm 0.0002$$

Power law decay with nontrivial exponent

Distribution of superior records

- Cumulative probability distribution $F_N(x)$ that:
 1. Sequence is superior ($X_n > A_n$ for all n) and
 2. Current record is larger than x ($X_N > x$)
- Gives the desired probability immediately

$$S_N = F_N(A_N)$$

- Recursion equation

$$F_{N+1}(x) = x F_N(x) + (1 - x) F_N(A_N) \quad x > A_{N+1}$$

old record holds a new record is set

- Recursive solution

$$F_1(x) = 1 - x$$

$$F_2(x) = \frac{1}{2} (1 + x - 2x^2)$$

$$F_3(x) = \frac{1}{18} (7 + 2x + 9x^2 - 18x^3)$$

$$F_4(x) = \frac{1}{576} (191 + 33x + 64x^2 + 288x^3 - 576x^4)$$

$$S_1 = \frac{1}{2}$$

$$S_2 = \frac{7}{18}$$

$$S_3 = \frac{191}{576}$$

$$S_4 = \frac{35393}{120000}$$

$$S_N = F_N(A_N)$$

\Rightarrow

Scaling Analysis

- Convert recursion equation

$$F_{N+1}(x) = x F_N(x) + (1 - x) F_N(A_N)$$

into a differential equation (N plays role of time!)

$$\frac{\partial F_N(x)}{\partial N} = (1 - x) [F_N(A_N) - F_N(x)]$$

- Seek a similarity solution ($N \rightarrow \infty$ limit)

$$F_N(x) \simeq S_N \Phi(s) \quad \text{with} \quad s = (1 - x)N$$

boundary conditions $\Phi(0) = 0$ and $\Phi(1) = 1$ $\left(1 - \frac{N}{N+1}\right) N \rightarrow 1$

- Similarity function obeys first-order ODE

$$\Phi'(s) + (1 - \beta s^{-1})\Phi(s) = 1$$

Similarity solution gives distribution of scaled record

Similarity Solution

- Equation with yet unknown exponent

$$\Phi'(s) + (1 - \beta s^{-1})\Phi(s) = 1$$

- General solution

$$\Phi(s) = s \int_0^1 dz z^{-\beta} e^{s(z-1)}$$

- Boundary condition dictates the exponent


$$\int_0^1 dz z^{-\beta} e^{(z-1)} = 1$$

- Root is a transcendental number

$$\beta = 0.450265027495$$

Analytic solution for distribution and exponent

Distribution of records



scaling variable $s = (1 - x)N$


Inferior records

- Start with sequence of random variables

$$\{x_1, x_2, x_3, \dots, x_N\}$$

- Calculate the sequence of records

$$\{X_1, X_2, X_3, \dots, X_N\} \quad \text{where} \quad X_n = \max(x_1, x_2, \dots, x_n)$$



- Compare with the expected average

$$\{A_1, A_2, A_3, \dots, A_N\} = \{1/2, 2/3, 3/4, \dots, N/(N+1)\}$$

- Inferior sequence = records always below average

$$X_n > A_n \quad \text{for all} \quad 1 \leq n \leq N$$

- What fraction of sequences are inferior?

$$I_N \sim N^{-\alpha}$$

expect power law decay, different exponent

Probability sequence is inferior

- Start with sequence of random variables

$$\{A_1, A_2, A_3, \dots, A_N\} = \{1/2, 2/3, 3/4, \dots, N/(N+1)\}$$

- One variable

$$x_1 < \frac{1}{2} \implies I_1 = \frac{1}{2}$$

- Two variables

$$x_1 < \frac{1}{2} \text{ and } x_2 < \frac{2}{3} \implies I_2 = \frac{1}{2} \times \frac{2}{3} = \frac{1}{3}$$

- Recursion equation (no interactions between variables)

$$I_{N+1} = I_N \frac{N}{N+1}$$

- Simple solution

$$I_N = \frac{1}{N+1} \quad I_N \sim N^{-1}$$

power law decay with trivial exponent

General distributions

- Arbitrary distribution function
- Single parameter contains information about tail

$$\alpha = \lim_{N \rightarrow \infty} N \int_{A_N}^{\infty} dx \rho(x)$$

- Equals the exponent for inferior sequences

$$I_N \sim N^{-\alpha}$$


- Exponent for superior sequences

$$\alpha \int_0^1 dz z^{-\beta} e^{\alpha(z-1)} = 1$$

- Power-law distributions (compact support)


$$R(x) \sim (1-x)^\mu \quad \Longrightarrow \quad \alpha = \left[\Gamma\left(1 + \frac{1}{\mu}\right) \right]^\mu$$

Continuously varying exponents



$$\alpha_{\min} \leq \alpha < \infty$$

$$\alpha_{\min} = e^{-\gamma} = 0.561459$$







$$\beta_{\max} = 0.621127$$

$$0 < \beta \leq \beta_{\max}$$

Tail of distribution function controls record statistics

Records in earthquake data inter-event times



good agreement with
theoretical predictions

Conclusions

- Studied persistent configuration of record sequences
- Linear evolution equations (but nonlocal/memory)
- Dynamic formulation: treat sequence length as time
- Similarity solutions for distribution of records
- Probability of persistent configuration (inferior, superior, inferior) decays as a power-law
- Power laws exponents are generally nontrivial
- Exponents can be obtained analytically
- Tail of distribution function controls record statistics